

external exposure, databanks on patterns of behaviors, and markers of internal exposure.

*The authors declare they have no actual or potential competing financial interests.*

**Martie van Tongeren**

**John W. Cherrie**

Institute of Occupational Medicine

Edinburgh, United Kingdom

E-mail: martie.vantongeren@iom-world.org

## REFERENCES

Lioy PJ, Rappaport SM. 2011. Exposure science and the exposome: an opportunity for coherence in the environmental health sciences [Editorial]. *Environ Health Perspect* 119:A466–A467.

Vineis P, Khan AE, Vlaanderen J, Vermeulen R. 2009. The impact of new research technologies on our understanding of environmental causes of disease: the concept of clinical vulnerability. *Environmental Health* 8:54; doi:10.1186/1476-069X-8-54 [Online 30 November 2009].

## An Integrated Approach to the Exposome: Rappaport and Lioy Respond

<http://dx.doi.org/10.1289/ehp.1104719R>

We welcome the remarks of van Tongeren and Cherrie regarding our recent editorial (Lioy and Rappaport 2011) and see no particular differences in our positions. As originally conceived, the exposome concept promoted investigations of disease etiology, that is, finding unknown causes of disease (Wild 2005). This requires an untargeted study design so that important, but as yet unrecognized, exposures will not be missed (Rappaport and Smith 2010). Such untargeted designs lend themselves to omic characterization of biospecimens (of the top-down type), as has been demonstrated in recent metabolomic investigations (e.g., Wang et al. 2011). Many external measurements of exposure focus on specific chemicals or classes of agents, but van Tongeren and Cherrie offer examples of untargeted designs (e.g., mining records of household food purchases). In any case, as measurements of external phenomena become less targeted, they become more exposomic (of the bottom-up type). The real issue is to recognize the underlying reasons for estimating exposure levels. If measurements are intended to find unknown sources of disease, then they are consistent with the exposome concept. If they are intended for other purposes (e.g., dose response, risk assessment/management, source characterization), then they follow more traditional lines of exposure assessment/science. As we emphasized in our editorial (Lioy and Rappaport 2011), both approaches have merit, and a combination of the two offers particular advantages for both identifying and preventing hazardous exposures, and thereby mitigating diseases.

*The authors declare they have no actual or potential competing financial interests.*

**Stephen M. Rappaport**

School of Public Health

University of California

Berkeley, California

E-mail: srappaport@berkeley.edu)

**Paul J. Lioy**

Environmental and Occupational Health

Sciences Institute

Robert Wood Johnson Medical School–

UMDNJ and Rutgers University

Piscataway, New Jersey

## REFERENCES

Lioy PJ, Rappaport SM. 2011. Exposure science and the exposome: an opportunity for coherence in the environmental health sciences [Editorial]. *Environ Health Perspect* 119:A466–A467.

Rappaport SM, Smith MT. 2010. Environment and disease risks. *Science* 330(6003):460–461.

Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, et al. 2011. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature* 472(7341):57–63.

Wild CP. 2005. Complementing the genome with an “exposome”: the outstanding challenge of environmental exposure measurement in molecular epidemiology. *Cancer Epidemiol Biomarkers Prev* 14(8):1847–1850.

## Application of the Ecohealth Model to Translate Knowledge into Action in the Health Sciences

<http://dx.doi.org/10.1289/ehp.1104847>

As noted by Barkin and Schlundt (2011), addressing the public health needs of the population using evidence from biomedical research necessarily requires a wholistic approach that is both multilevel and multidisciplinary. Although there may be public health benefits, there are also important challenges when generating knowledge, at the microenvironmental level, as well as at the macroenvironmental level. This happens particularly when evidence is translated into interventions that generate benefits for all who are involved in the health process; for example, in dealing with obesity, these interventions would benefit users, the health system, food producers, and others. To complement the response to these challenges, we suggest a greater application of the ecohealth model. This model has been proposed as a new analytical model for research action based on the ecosystemic approach to human health, an approach that places health within the realm of the environment and acknowledges cause–effect interconnections between human health and humans’ biophysical, social, and economic environment.

The ecohealth model stems from the generation of health knowledge and the multiple interconnections between the different components of the ecosystem. It sets forth that

these interconnections are complex and interdependent and include social determinants and disparities, as well as biophysical determinants. From this perspective, scientists need to revise their models and research methods and open up to new analytical focuses and new forms of collaboration and interaction, going beyond the biophysical characteristics of systems and the scientific community itself. For many reasons, the traditional methods used in the study of the micro–macro environment have not been able to fulfill the expectations for health and welfare or those for improving sanitary conditions of populations. Thus, we need to periodically evaluate evaluations and adjust programs, interventions, and health policies.

Although traditional methods take into account the economy and the community, often at the expense of the environment (jeopardizing the possibility of a sustainable ecosystem), the ecohealth model breaks up each of its components into different categories (Hancock 1990; Lebel 2005). It confers equal importance to environmental management, economic factors, and the community’s aspirations, and it places human health at the center of the intersection of these three elements. In this sense, the ecohealth model itself is part of the sustainable development process, and its fundamental premise is to be inclusive. Interventions and health programs based on evidence generated under the ecohealth model should be more cost-effective than many medical treatments or traditional healthcare interventions. This analytical model and its methodological research approach involve three participating groups: researchers and other specialists; community members, such as common citizens, businessmen, farmers, fishermen, and miners; and decision makers in health interventions. Besides the need for the participation of these three groups, the ecohealth model is based on three methodological pillars: transdisciplinarity, participation and equity.

- Transdisciplinarity implies a multilevel and translevel vision, with a broad scope and collaboration in the study of health determinants and conditions related to the ecosystem.
- Participation intends to achieve consensus on the definition of the study’s objective among scientists, community members, and decision makers, both between and within groups.
- Equity includes the analysis of the roles of men and women and their different degrees of influence in decisions on access to and use of financial resources, as well as equity in benefits and rewards for all of those involved in a concrete health problem.

Each of these pillars generates, to a great extent, conditions for a more effective and

efficient translation of scientific knowledge into action, as well as addressing the challenges set forth by Barkin and Schlundt (2011). In fact, the transdisciplinarity component responds to the need for greater collaboration between researchers from all involved disciplines, as well as other social actors studying the same problem. Participation and equity in the involved groups (researchers, decision makers, and community members) at the time of implementing interventions and generating benefits would guarantee a greater effectiveness in the design and implementation of interventions with an ecohealth focus. These components would also promote greater equality and equity in the benefits and rewards for all involved parties.

In summary, the collaboration between researchers during generation and application of scientific evidence—at both the microenvironmental and macroenvironmental levels—can guarantee greater benefits, acceptability, and effectiveness in interventions at the population level for all of those involved.

*The authors declare they have no actual or potential competing financial interests.*

**Armando Arredondo  
Emanuel Orozco**

Center for Health Systems Research  
National Institute of Public Health  
Cuernavaca, Morelos, Mexico  
E-mail: armando.arredondo@insp.mx


#### REFERENCES

- Barkin S, Schlundt D. 2011. The challenge facing translation of basic science into clinical and community settings to improve health outcomes [Editorial]. *Environ Health Perspect* 119:A418–A419.
- Hancock T. 1990. Toward Healthy and Sustainable Communities: Health, Environment and Economy at the Local Level. A Presentation at the 3rd Symposium on Environmental Health, Quebec, November 22, 1990. Toronto, Ontario, Canada: Faculty of Environmental Studies, York University.
- Lebel J. 2005. Salud un Enfoque Ecosistemático [in Spanish]. Bogotá, Colombia: EnFoco - Alfaomega/International Development Research Centre.

# Stay Connected

This iPhone app provides easy access to

- » Ahead-of-Print research articles, reviews, and commentaries
- » Current issue
- » Archived issues
- » *The Researcher's Perspective* podcast series



Get it for your iPhone or iPod Touch. Get it for FREE.



Scan with your  
iPhone or iPod Touch  
QR reader  
to get the *EHP* app.

To download the free app  
visit the iTunes App Store, or  
log on to  
[www.ehponline.org/mobile](http://www.ehponline.org/mobile)